

**Roger Williams University**  
**DOCS@RWU**

---

Feinstein College of Arts & Sciences Faculty  
Publications

Feinstein College of Arts and Sciences

---

2005

# A fossil reef from the last interglacial, Western Australia

Benjamin J. Greenstein  
*Roger Williams University, [bgreenstein@rwu.edu](mailto:bgreenstein@rwu.edu)*

John M. Pandolfi

David R. Blakeway

Follow this and additional works at: [https://docs.rwu.edu/fcas\\_fp](https://docs.rwu.edu/fcas_fp)




Part of the [Biology Commons](#), and the [Marine Biology Commons](#)

---


## Recommended Citation

Greenstein, Benjamin. J., John M. Pandolfi, and David R. Blakeway. "A fossil reef from the Last Interglacial, Western Australia." *Coral Reefs*, v. 24, no. 4, 2005, pp. 593.

This Article is brought to you for free and open access by the Feinstein College of Arts and Sciences at DOCS@RWU. It has been accepted for inclusion in Feinstein College of Arts & Sciences Faculty Publications by an authorized administrator of DOCS@RWU. For more information, please contact [mwu@rwu.edu](mailto:mwu@rwu.edu).



**Fig. 1** Location of the Sandalwood Peninsula



**Fig. 2** Fossil coral reef assemblage. **a** Coral colonies including *Goniastrea*, *Hydnophora*, *Favia*, *Favites*, *Lobophyllia* and *Platygyra* preserved in growth position on erosion surface. **b** Detail of erosion surface with *Goniastrea australiensis* preserved in situ. **c** View of erosion surface with extensive borings preserved

## A fossil reef from the last interglacial, Western Australia

Received: 29 June 2005

Accepted: 16 August 2005

Published online: 6 October 2005

© Springer-Verlag 2005

We report an initial investigation of a Devils Point Event (DPE) style surface exposed approximately 1.5 m above present sea level in a spectacularly preserved reef sequence in southern Exmouth Gulf, Western Australia. The fossil coral reef is exposed at Point Maxwell on the Sandalwood Peninsula (Fig. 1). Two reef-building sequences, separated by a distinctive erosion surface are present. The erosion surface truncates a lower coral-bearing unit and has been colonized by numerous coral colonies preserved in growth position (Fig. 2a, b). The surface is riddled with many different borings (Fig. 2c). Although initial dating of corals using alpha spectrometry suggests a Late Pleistocene age, the geochronology of the sequence has not been established. We interpret it as Last Interglacial (MIS 5e) based on its position relative to present sea level, its mode of preservation, and a tentative correlation with fossil reefs exposed on the west side of Cape Range (for which radiometric ages have been published). The distinctive sedimentologic and paleontologic signatures of the erosion surface suggest a regressive-transgressive event similar to the DPE described from Great Inagua and San Salvador Islands, Bahamas by Wilson et al. (1998) and White et al. (1998). U-series dates obtained from reef corals preserved in the Pleistocene Tantabiddi Limestone exposed on the western margin of Cape Range (approximately 75 km west of southern Exmouth Gulf) were published by Stirling et al. (1998) and evaluated in the context of the DPE by White et al. (2001). Reliable dates for that region range from 115 to 129 ka, and are divided equally between pre- and post-DPE ages. However, the stratigraphic relationship between the assemblages from which the corals were obtained is unknown. Moreover, the significance of Late Cenozoic tectonism versus glacio-eustatic change in exposing the marine sequences in the region remains unclear (Kendrick et al. 1991). Further examination of the fossil reef described here should provide additional evidence to assess the climatic instability that has been reported for the Last Interglacial.

**Acknowledgements** We thank Mark Wilson (The College of Wooster, USA) for feedback on the borings we observed.

### References

- Kendrick GW, Wyrwoll K-H, Szabo BJ (1991) Pliocene-Pleistocene coastal events and history along the western margin of Australia. *Quaternary Sci Rev* 10:419–439
- Stirling CH, Esat TM, Lambeck K, McCulloch MT (1998) Timing and duration of the Last Interglacial: evidence for a restricted interval of widespread coral reef growth. *Earth Planet Sc Lett* 160:745–762
- White B, Curran HA, Wilson MA (1998) Bahamian coral reefs yield evidence of a brief global sea-level lowstand during the Last Interglacial. *Carbonates and Evaporites* 13:10–22
- White B, Curran HA, Wilson MA (2001) A sea-level lowstand (Devil's Point Event) recorded in Bahamian reefs: comparison with other Last Interglacial climate proxies. In: Greenstein BJ, Carney CK (eds) *Proceedings of the 10th symposium on the geology of the Bahamas and other carbonate regions*. Gerace Research Center, San Salvador, Bahamas, pp 109–128
- Wilson MA, Curran HA, White B (1998) Paleontological evidence of a brief global sea-level event during the last interglacial. *Lethaia* 31:241–250

B. J. Greenstein (✉) · J. M. Pandolfi · D. R. Blakeway  
Department of Geology, Cornell College, Mt. Vernon IA 52314, USA  
E-mail: bgreenstein@cornellcollege.edu  
Tel.: +1-319-8959845  
Fax: +1-319-8955667

Centre for Marine Studies and Department of Earth Sciences, University of Queensland, Brisbane 4072 QLD, Australia  
School of Earth and Geographical Sciences, University of Western Australia, Crawley 6009, Australia

# Reef sites

Coral Reefs (2005) 24: 593  
DOI 10.1007/s00338-005-0043-z